# NEXT GENERATION & NEXT PARADIGM

# NEXPHARM KOREA



The 30th Anniversary 1995~2025

NEXPHARM KOREA

To the world,
For your healthier tomorrow,

# NEXPHARM KOREA

Based on remarkable technologies,
We manufacture & provide

## the best quality medicines



# 4
## DOSAGE FORMS

- TABLETS
- CAPSULES
- GRANULES & POWDERS
- LIQUIDS(SOLUTION & SUSPENSION)

# 3
## FACTORIES

PIC/S GMP
COMPLIANCE

- NEXPHARM KOREA Osong Plant
- NEXPHARM KOREA Bio Plant
- EDEN PHARMA Plant (subsidiary)




# Research Performances

**10**
**PATENTS**



**41**
**CLINICALS**



**Technology Transfer**



# R&D Pipe Lines

| Classification | Project | Indication | Research | Non-clinical study | Clinical study | | |
|---|---|---|---|---|---|---|---|
| | | | | | Phase 1 | Phase 2 | Phase 3 |
| Incrementally modified drugs | NEX004 | Peptic ulcer | | | | | |
| New natural substances | G47 Extract | Hyperlipidemia | | | | | |
| | G46 Extract | Osteoarthritis | | | | | |
| Biopharmaceutical new drug | TRAP80 | Arteriosclerosis | | | | | |
| Biobetter | NEX005 | Diabetes | | | | | |
| Individually recognized functional ingredient | NEX006 | Macular degeneration | | | | | |
| | NEX006 compound | Macular degeneration | | | | | |
| | NEX007 | Triglycerides | | | | | |

# PRODUCT LIST

## NEW DRUGS

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) | STATUS |
|---|---|---|---|---|---|
| **NICER TAB. 30MG** |  | Nicergoline 30mg | Treatment of senile dementia, migraines of vascular origin, transient ischemia, platelet hyper-aggregability, and macular degeneration. | 1 tablet | Launched |
| **MEOLIGA SYRUP** |  | 1 sachet (10ml) contains, Caffeine Anhydrous 16.67mg Dimenhydrinate 33.33mg Pyridoxine hydrochloride 4.17mg | Prevention and relief of dizziness, vomiting, and headache caused by motion sickness. | 3-4 sachets | Launched |
| **MILFRAN CAP. 50MG** | | Milnacipran 50mg | Treatment of fibromyalgia, depression | 1 capsule, two times | 2026 |
| **PIESTAR TAB. 2/10 & 4/10** | | Pitavastatin 2mg/ Ezetimibe 10mg & Pitavastatin 4mg/ Ezetimibe 10mg | Primary hypercholesterolemia | 1 tablet | 2027 |
| **EZEFEN TAB. 10/160** | | Fenofibrate 160mg Ezetimibe 10mg | Adjunct diet to reduce elevated total cholesterol (total-C), low-density lipoprotein cholesterol (LDL-C), apolipoprotein B (Apo B), and non-high-density lipoprotein cholesterol (non-HDL-C) in patients with mixed hyperlipidemia | 1 tablet | 2028 |

# PRODUCT LIST

## GASTROINTESTINAL SYSTEM

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| **LAFUDILE TAB.** |  | Lafutidine 10mg | **H2-Blocker**<br>1. Gastric and duodenal ulcers<br>2. Gastric mucosal lesions of acute and chronic gastritis<br>3. Helicobacter pylori infected peptic ulcer<br>4. Pre-anesthetic medication<br>5. Reflux esophagitis | 1 tablet,<br>1 ~ 2 times |
| **NITADINE TAB.** |  | Nizatidine 75mg | **H2-Blocker**<br>Hyperacidity, Heartburn | 1 tablet,<br>2 times |
| **ESORA TAB. 20MG** |  | Esomeprazol magnesium dihydrate 21.69mg(20mg as esomeprazole) | **PPI**<br>1. Short-term treatment of heartburn and other symptoms of gastroeso- phageal reflux disease (GERD)<br>2. Short-term treatment in healing and symptomatic resolution of erosive esophagitis<br>3. Maintenance of symptom resolution and healing of erosive esophagitis<br>4. Combination with amoxicillin and clarithromycin for treatment of Helicobacter pylori infection and duodenal ulcer to eradicate H. pylori<br>5. Reduction in occurrence of gastric ulcers associated with continuous NSAID therapy in patients at risk of developing gastric ulcers | 1 tablet,<br>2 times |
| **ESORA TAB. 40MG** |  | Esomeprazol magnesium dihydrate 43.38mg(40mg as esomeprazole) | | 1 tablet,<br>once a day |
| **NEXONA TAB. 2MG** |  | Irsogladine maleate 2mg | **Antiulcerants**<br>1. Gastric ulcer<br>2. Gastric mucosal lesions (erosion, bleeding, redness & edema) in acute gastritis & acute exacerbation of chronic gastritis | 4mg,<br>1 ~ 2 in divided dose |
| **NEXONA TAB. 4MG** |  | Irsogladine maleate 4mg | | |
| **MUCOBA TAB.** |  | Rebamipide 100mg | **Antiulcerants**<br>1. Gastric ulcers.<br>2. Gastric mucosal lesions (erosion, bleeding, redness & edema) in acute gastritis & acute exacerbation of chronic gastritis | 1 tablet,<br>3 times |
| **NEXTIL TAB.** |  | Artemisia asiatica 95% ethanol extract(20→1) 60mg | **Antiulcerants**<br>1. Gastric mucosal lesions (acute & choronic gastritis)<br>2. Prevention of gastritis caused by NSAIDs administration | 1 tablet,<br>3 times |
| **NEXIGEL SUS.** |  | Oxethazaine 20mg<br>Dried aluminum hydroxide gel 582mg<br>Magnesium hydroxide 196mg/10mL | **Antacids**<br>Acute and chronic pain, gastritis, duodenal ulcer, esophagitis, gastric pain caused by digestive disorders during radiation therapy | 1 stick,<br>4 times |
| **NEXTO TAB.** |  | Itopride hydrochloride 50mg | **GI tract regulators & antispasmodics**<br>GI symptoms associated with func- tional dyspepsia i.e. sensation of bloating, early satiety, upper abdominal pain or discomfort, anorexia, heartburn, nausea & vomiting | 1 tablet,<br>3 times |

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| **SOLCON TAB.** |  | Polycarbophil calcium 625mg | **Laxatives**<br>Chronic constipation, non-specific diarrhea | 2 tablets,<br>1 ~ 4 times |
| **NEXCOLON TAB.** |  | Monobasic sodium phosphate monohydrate 1,102mg<br>Dibasic sodium phosphate anhydrous 398mg | Clean out of intestine before colonos- copy | Evening before colonoscopy<br>: 4 tablets,<br>5 times in every 15 minutes<br>Colonoscopy day : 4 tablets<br>3 ~ 5 hours before,<br>3 times in every 15 minutes |

# ANTIHYPERLIPIDEMIAS

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| **APITO TAB. 10MG** |  | Atorvastatin calcium trihydrate 10.85mg | **Antilipidemic agents**<br>Reduction of elevated cholesterol & triglyceride levels if other measures eg diet, exercise, wt reduction have been unsuccessful in patients with primary hypercholesterolaemia or mixed hyperlipidaemia (Fredrickson types IIa & IIb), hereditary familial heterozygous or homozygous hypercholesterolaemia | 1 tablet,<br>1 ~ 2 times |
| **APITO TAB. 20MG** |  | Atorvastatin calcium trihydrate 21.7mg | | 1 tablet,<br>once a day |
| **ROSUMIN TAB. 5MG** |  | Rosuvastatin calcium 5.2mg | **Antilipidemic agents**<br>1. Primary hyperlipidemia or Mixed dyslipidemia (type IIb)<br>2. Homozygous familial hypercholesterolemia<br>3. Reducing the risk of atherosclerosis by lowering total cholesterol levels and LDL(low-density lipoprotein) cholesterol levels.<br>4. Heterozygous familial hypercholesterolemia<br>5. Primary dysbetalipoproteinemia<br>6. Reducing the risk of cardiovascular diseases | 1 tablet,<br>once a day |
| **ROSUMIN TAB. 10MG** |  | Rosuvastatin calcium 10.4mg | | |
| **ROSUMIN TAB. 20MG** |  | Rosuvastatin calcium 20.8mg | | |
| **EZETIN TAB.** |  | Ezetimibe 10mg | **Antilipidemic agents**<br>1. Primary hypercholesterolemia (heterozygous familial and nonfamilial)<br>2. Homozygous familial hypercholesterolemia<br>3. Homozygous Sitosterolemia (Plantosterolemia) | 1 tablet,<br>once a day |

# PRODUCT LIST

## CARDIOVASCULAR SYSTEM

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| KARYPT GRANULE |  | Polystyrene sulfonate calcium 5g | Hyperkalemia | 15 ~ 30g in 2 ~ 3 divided doses |
| OMAPRON TAB. |  | Limaprost alpha-cyclodextrin (alfadex) 166.67μg (5μg as Limaprost) | **Vasodilators**<br>1. Improvement of various ischemic symptoms such as ulcer , pain and feeling of coldness associated with obliterative thromboangitis (Buerger's disease)<br>2. Improvement of subjective symptoms (pain and numbness of lower legs) and gait ability associated with acquired lumbar spinal canal stenosis (in patients with bilateral intermittent claudication showing normal SLR test result) | 1 ~ 2 tablets, 3 times |

## ANTIDIABETES

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| FLOGA TAB. 5MG |  | Dapagliflozin propanediol monohydrate 6.15mg | **SGLT-2 inhibitor**<br>Type 2 Diabetes: This drug is administered as an adjunct to diet and exercise to improve blood sugar control in patients with type 2 diabetes.<br>- Monotherapy<br>- Combination therapy | 1 tablet, once a day |
| FLOGA TAB. 10MG |  | Dapagliflozin Propanediol Monohydrate 12.30mg | | 1 tablet, once a day |
| DIMERYL TAB. 2MG |  | Glimepiride 2mg | **Sulfonylureas**<br>Type 2 diabetes mellitus | 1 ~ 4mg, once a day before breakfast, up to 8mg a day |
| ACTORIN TAB. 15MG |  | Pioglitazone hydrochloride 16.53mg | **Thiazolidinediones**<br>Type 2 diabetes mellitus | 1 ~ 2 tablets, once a day |
| ACTORIN TAB. 30MG |  | Pioglitazone hydrochloride 33.06mg | **Thiazolidinediones**<br>Type 2 diabetes mellitus | 1 tablets once a day |

## ANTIDIURETICS

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| NURITAN TAB. 0.1MG |  | Desmopressin acetate 0.1mg | **Pituitary hormone**<br>1. Primary nocturnal enuresis (over 5 years old)<br>2. Treatment of nocturia symptom related to nocturnal polyuria (Adults only) | 1 ~ 2 tablets, once a day |
| NURITAN TAB. 0.2MG |  | Desmopressin acetate 0.2mg | | 1 tablet, once a day |

# LIVER TONICS

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| HEPATO CAP. |   | Ursodeoxycholic acid 50mg Biphenyl Dimethyl Dicarboxylate 12.5mg | **Hepatic protectors** Chronic liver disease with elevated SGPT | 1 capsule, 3 times |

# CENTRAL NERVOUS SYSTEM

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| NEXTIMIN CAP. 1.5MG |   | Rivastigmine tartrate 2.4mg | **Nootropics** Dementia of the Alzheimer's type Dementia of the Parkinson's disease | 1 capsule, 2 times |
| NEXTIMIN CAP. 3MG |   | Rivastigmine tartrate 4.8mg | | 1 capsule, 2 times |

# ANTIDIPRESSION DRUGS

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| DESVELA TAB. 50MG |   | Desvenlafaxine benzoate 73.185mg | **Antidipression** Depression | 1 tablet, once a day |
| DESVELA TAB. 100MG |   | Desvenlafaxine benzoate 146.37mg | | 1 tablet, once a day |

# UROLOGICAL SYSTEM

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| NEOTRIN TAB. |   | Terazosin hydrochloride 2.374mg | Dysuria due to benign prostatic hyperplasia Hypertension (mild to moderate) | 1 tablet, once a day |

# PRODUCT LIST

## ANTIBIOTICS

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| **FOSFO POWDER** |  | Fosfomycin trometamol 5.631g (3g as fosfomycin) | **Antibiotic**<br>1. Acute symple cystitis in adult women<br>2. Severe asymptomatic bacteriuria in pregnanacy<br>3. Prevention of infection before and after transrectal prostate biopsy in adult men | 1 stick, once a day |
| **LEFOXACIN TAB.** |  | Levofloxacin hydrate 100mg | **Quinolones**<br>Mild, moderate and severe infections caused by susceptible strains of the designated microorganisms for the following conditions: Tablet (250,500 mg) and Solution for Infusion (500 mg): Acute bacterial sinusitis, acute bacterial exacerbation of chronic bronchitis, nosocomial pneumonia, community- acquired pneumonia, complicated skin and skin structure infections, uncomplicated skin and skin structure infections, chronic bacterial prostatitis, complicated urinary tract infections, acute pyelonephritis and uncomplicated urinary tract infections. Tablet (750 mg): Acute bacterial sinusitis, complicated skin and skin structure infections, community-acquired pneumonia and nosocomial pneumonia. | 1 tablet, 2 ~ 3 times |

## ANTIINFECTIVES

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| **IDIZOLE CAP.** |  | Itraconazole pellet 454.55mg (100mg as Itraconazole) | **Antiinfection**<br>Antifungal agent such as : Candida vaginitis, Ginger, Ringworm of the body caused by dermatophytes, ringworm on the upper part of the body, ringworm on the hands, ringworm on the feet, Oral candidiasis, Fungal keratitis, Onychomycosis, Systemic fungal infections | 1 ~ 2 capsules, 1 ~ 2 times |
| **DICAZOLE CAP.** |  | Fluconazole 50mg | **Antifungal agent**<br>Cryptococcosis, generalised candi- diasis, mucosal candidiasis including oropharyngeal & vag candidiasis.<br>Prevention of fungal infections in immunocompromised patients | Onychom- ycosis : 150mg, once a week |
| **FAMCILON TAB.** |  | Famciclovir 250mg | **Antiviral agent**<br>Treatment of herpes zoster (shingles) in immunocompet ent patients ; treatment or suppression of recurrent genital herpes in immunocompetent patients; treatment of recurrent episodes of orolabial or genital herpes in HIV-infected adults; treatment of recurrent herpes labialis (cold sores) in immunocompetent patients | 125 ~ 250mg, 2 ~ 3 times |

# RESPIRATORY SYSTEM

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| LEVRONE TAB. |  | Levodropropizine 60mg | Cough due to acute and chronic bronchitis | 1 tablet, 3 times |

# NSAIDs

| BRAND NAME | | COMPOSITION | INDICATIONS | DOSAGE (/DAY) |
|---|---|---|---|---|
| DEXTALONE TAB. |  | Dexibuprofen D.C. 400mg | 1. Rheumatoid arthritis<br>2. Arthrosis<br>3. Ankylosing spondylitis<br>4. Pain or inflammation after surgery, injury<br>5. Supportive treatment for infection accompanying inflammation, pain, fever | 1 tablet, 2 ~ 3 times |
| ASNAC TAB. |  | Aceclofenac 100mg | Rheumatoid arthritis, ankylosing spondylitis, osteoarthritis, pain and inflammation associated with musculoskeletal and joint disorders, low back pain | 1 tablet, 2 times |
| CENEX CAP. 100MG |  | Celecoxib 100mg | **COX-2 Selective NSAID**<br>1. Osteoarthritis<br>2. Rheumatoid arthritis<br>3. Ankylosing spondylitis<br>4. Acute pain : after surgery, tooth extraction<br>5. Primary dysmenorrhea | 1 capsule, 2 times |
| CENEX CAP. 200MG |  | Celecoxib 200mg | | 1 capsule, once a day |
| CENEX CAP. 400MG |  | Celecoxib 400mg | | 1 capsule, once a day |
| THERAPEN TAB. |  | Acetaminophen 325mg<br>Tramadol hydrochloride 37.5mg | Moderate to severe acute & chronic pain | 2 tablets, 4 times |
| THERAPEN- SEMI TAB. |  | Acetaminophen 162.5mg<br>Tramadol hydrochloride 18.75mg | Moderate to severe acute & chronic pain | Initial : 4 tablets, and then, every 6 hours, up to 16 tablets |

Designed by [ (주)비비아이커뮤니케이션즈 02 2264 8355 ]

넥스팜코리아
NEXPHARM KOREA

**HEAD Office**
KD building, 36, Yangjaecheonro 19-gil Seocho-gu, Seoul, Repulbic of Korea
TEL: 02-3473-6365 FAX: 02-3473-6366

**Factory**
168-41, Osongsaengmyeong 4-ro, Osong-eup, Heungdeok-gu, Cheongju-si, Chungbuk, Republic of Korea
TEL: 043-716-2080 FAX: 070-4649-5339

http://www.nexpharm.co.kr

EDEN PHARMA

**HEAD Office**
KD building, 36, Yangjaecheonro 19-gil Seocho-gu, Seoul, Repulbic of Korea
TEL: 02-3487-7005 FAX: 02-6952-1660

**Factory**
1149, Nongdari-ro, Munbaek-myeon, Jincheon-gun, Chungbuk, Republic of Korea
TEL: 043-534-9100 FAX: 070-8220-7610

http://www.edenpharma.co.kr